

# QUALI-PRO

GROUP 3 | 11 FUNGICIDE

ACCEPTED  
FOR REGISTRATION

March 1, 2018

DOC ID 555792

New York State Department  
of Environmental Conservation  
Division of Materials Management  
Pesticide Product Registration



# STROBE® PRO

## BROAD SPECTRUM FUNGICIDE

For control of Turf Diseases.

ACTIVE INGREDIENTS:	% BY WT.
Azoxystrobin*	6.0%
Propiconazole**	10.0%
OTHER INGREDIENTS:	84.0%
TOTAL:	100.0%

A liquid soluble concentrate that contains 0.87 lbs. propiconazole and 0.52 lbs. azoxystrobin per gallon.

\*IUPAC: methyl (E)-2-[2-[6-(2-cyanophenoxy)pyrimidin-4-yloxy]phenyl]-3-methoxyacrylate

\*\*CAS Number 60207-90-1

KEEP OUT OF REACH OF CHILDREN  
**CAUTION**

EPA Reg. No. 53883-418

EPA Est No. 53883-TX-002

## CONTENTS: 1 GALLON



0 72693 40076 4  
EPA 111417



Manufactured for:

## Control Solutions Inc.

A member of Adama  
Consumer and Professional Solutions  
5903 Genoa-Red Bluff, Pasadena, TX 77507

KEEP OUT OF REACH OF CHILDREN

## CAUTION

FIRST AID	
<b>IF SWALLOWED:</b>	<ul style="list-style-type: none"><li>• Call a poison control center or doctor immediately for treatment advice.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• Do not induce vomiting unless told to by the poison control center or doctor.</li><li>• Do not give anything by mouth to an unconscious person.</li></ul>
<b>IF INHALED:</b>	<ul style="list-style-type: none"><li>• Move person to fresh air.</li><li>• If person is not breathing, call 911 or an ambulance; then give artificial respiration, preferably by mouth-to-mouth if possible.</li><li>• Call a poison control center or doctor for further treatment advice.</li></ul>
<b>IF IN EYES:</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes; then continue rinsing.</li><li>• Call a poison control center or doctor for treatment advice.:</li></ul>
<b>IF ON SKIN:</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact SafetyCall® International for emergency medical treatment at (866) 897-8050.	

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Harmful if swallowed. Harmful if inhaled. Causes moderate eye irritation. Avoid breathing spray mist. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

#### PERSONAL PROTECTIVE EQUIPMENT (PPE)

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separate from other laundry. Some materials that are chemically resistant to this product are listed below.

#### Applicators and other handlers must wear:

1. Long-sleeved shirt and long pants
2. Protective eyewear
3. Chemical-resistant gloves made of any waterproof material such as polyvinyl chloride, nitrile rubber or butyl rubber
4. Shoes plus socks

## **USER SAFETY RECOMMENDATIONS**

### **Users Should:**

- 1.) Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- 2.) Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

## **ENVIRONMENTAL HAZARDS**

Azoxystrobin, can be persistent for several months or longer. Azoxystrobin has degradation products which have properties similar to chemicals which are known to leach through soil to ground water under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow may result in ground water contamination.

Azoxystrobin and propiconazole are toxic to freshwater and estuarine/marine fish;propiconazole is toxic to fish and shrimp and azoxystrobin is toxic to aquatic invertebrates. Do not apply directly to water. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

Notify state and/or Federal authorities and Control Solutions immediately if you observe any adverse environmental effects due to use of this product.

## **PHYSICAL OR CHEMICAL HAZARDS**

Do not mix or allow coming in contact with oxidizing agents. Hazardous chemical reaction may occur.

# **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements, specific to your State or Tribe, consult the agency responsible for pesticide regulation.

# **PRECAUTION**

**FAILURE TO FOLLOW THE USE DIRECTIONS AND RESTRICTIONS ON THIS LABEL MAY RESULT IN PLANT INJURY OR POOR DISEASE CONTROL.**

## AGRICULTURAL USES

For use to control diseases on turf on (Sod Farms)

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions, and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Exception: if product is soil injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is:

1. Coveralls
2. Chemical-resistant gloves made of any waterproof material such as polyvinyl chloride, nitrile rubber or butyl rubber
3. Shoes plus socks
4. Protective eyewear (goggles, face shield or safety glasses)

## NON AGRICULTURAL USES

For use to control diseases on turf on golf courses, lawns and landscape areas around residential, institutional, public, commercial and industrial buildings, parks, recreational areas and athletic fields.

### NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are **NOT** within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. The area being treated must be vacated by unprotected persons.

Do not treat areas while unprotected humans or domestic animals are present in the treatment areas.

Because certain states may require more restrictive reentry intervals, consult your State Department of Agriculture for further information.

Do not allow entry into treatment area until area that was treated with Strobe® Pro is dry.

## PRODUCT INFORMATION

Strobe® Pro is a combination of two broad spectrum, preventative fungicides with systemic properties for the control of many important turfgrass diseases. Strobe® Pro may be applied as a foliar spray in alternating spray programs or in tank mixes with other registered, turf protection products. All applications must be made according to the use directions that follow.

### PRODUCT USE RESTRICTIONS

- Do not graze or feed clippings from treated turf areas to animals.
- Do not apply Strobe® Pro to turf by air.
- Chemigation - Do not apply this product through any irrigation/chemigation system.

## PRODUCT USE PRECAUTIONS

- Strobe® Pro has demonstrated some phytotoxic effects when mixed with products that are formulated as EC's. These effects are enhanced if applications are made under cool, cloudy conditions and these conditions remain for several days following application. In addition, adjuvants that contain some form of silicone have also contributed to phytotoxicity.
- Bermudagrass can be sensitive to Strobe® Pro.

## INTEGRATED PEST (DISEASE) MANAGEMENT

Strobe® Pro should be integrated into an overall disease and pest management strategy whenever the use of a fungicide is required. Cultural practices known to reduce disease development should be followed. The **SPECIFIC USE DIRECTIONS** section in this label identifies specific IPM recommendations. Consult your local turf authority for additional IPM strategies established for your area. Strobe® Pro may be used in State Agricultural Extension advisory (disease forecasting) programs which recommend application timing based on environmental factors favorable for disease development.

## RESISTANCE MANAGEMENT

Strobe® Pro is a mixture of a Group 3 (propiconazole) and a Group 11 (azoxystrobin) fungicide. Strobe® Pro has two modes of action: (1) DMI (Demethylation Inhibitor of sterol biosynthesis) [Group 3], and (2) inhibitor of the Qo (quinone outside) site within the electron transport system (Qoi) as well as disrupting membrane synthesis by blocking demethylation [Group 11]. Fungal pathogens can develop resistance to products with the same mode of action when used repeatedly. Strobe® Pro is not cross-resistant with other classes of fungicides which have different modes of action. Because resistance development cannot be predicted, use of this product must conform to resistance management strategies established for turf in your area. Consult your local or State turf authorities for resistance management strategies that are complementary to those in this label.

Resistance management strategies may include rotating and/or tank-mixing with products having different modes of action or limiting the total number of applications per season. Avoid alternation with other Qo's, including pyraclostrobin and trifloxystrobin or other DMI's including triadimefon, fenarimol, and myclobutanil. Strobe® Pro must not be alternated or tank mixed with fungicides to which resistance has already developed. Continual use of A Strobe® Pro may allow less sensitive strains of pathogens to increase in the population and reduce the efficacy of Strobe® Pro. Albaugh encourages responsible resistance management to ensure effective long-term control of the fungal diseases on this label.

Use of a multiple-site contact fungicide including Chlorothalonil as an alternation or tank-mix partner can help alleviate selection pressure on pathogens and delay the onset of resistance. However, **DO NOT** use products containing chlorothalonil on home lawns or turf sites associated with apartment buildings, daycare centers, playgrounds, playfields, recreational park athletic fields, athletic fields located on or next to schools (i.e., elementary, middle and high schools), campgrounds, churches and theme parks.

Since pathogens differ in their potential to develop resistance to fungicides, the **SPECIFIC USE DIRECTIONS** section in this label provides resistance management strategies specific for each disease.

## SPRAYING/MIXING INSTRUCTIONS

### Restriction

Do not apply Strobe® Pro to turf by air. Do not exceed 3 fl. oz./1,000 sq. ft. every 30 days on any variety of bermudagrass. In FL, Do not apply Strobe® Pro to bermudagrass golf course greens when temperatures exceed 90°F.

Strobe® Pro may be applied with all types of spray equipment commonly used for making ground applications. Do not apply Strobe® Pro through any type of ultra-low volume (ULV) spray system (less than 3 gals/A). Proper adjustments and calibration of spraying equipment to give good canopy penetration and coverage is essential for good disease control. The higher rates in the rate range and/or shorter spray intervals may be required under conditions of heavy infection pressure, highly susceptible varieties, or when disease conducive environmental conditions exist

Apply Strobe® Pro in sufficient water volume for adequate coverage.

### SPRAY DRIFT MANAGEMENT

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, relative humidity) and method of application can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product. Do not apply at wind speeds greater than 15 mph. Apply as a medium or coarser spray (ASAE Standard 572). If applying at wind speeds less than 3 mph, the applicator must determine if a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

Applicators must follow all state and local pesticide drift requirements regarding application of propiconazole. Where states have more stringent regulations, they must be observed. All application equipment must be properly maintained and calibrated using appropriate carriers or surrogates.

### ATTENTION

**DO NOT** spray when conditions favor drift beyond area intended for application. Conditions which may contribute to drift include thermal inversion, wind speed and direction, sprayer nozzle/pressure combinations, spray droplet size, etc. Contact your State extension agent for spray drift prevention guidelines in your area.

### AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR

Do not apply when weather conditions favor drift from treated areas to non-target aquatic habitat.

### MIXING INSTRUCTIONS

To prepare spray solution, partially fill the spray tank with clean water and begin agitation. Add the specified amount of Strobe® Pro to the tank, allowing time for good dispersion, then add an adjuvant, if specified. If tank mixes are required, product must be added to the spray tank in the following order: Strobe® Pro, wettable granular or dry flowable formulations, wettable powders and flowable (aqueous suspensions) products. Finish filling the tank to the desired volume to obtain the proper spray concentration. Maintain agitation throughout the spraying operation.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Do not allow spray mixture to stand overnight or for prolonged periods. Make up only the amount of spray required for immediate use. Sprayers must be thoroughly cleaned immediately after application.

Do not use silicone-based products with Strobe® Pro due to possible phytotoxicity.

If spray-tank mixture is unsprayed for more than 18 hours (overnight), re-suspend product with agitation for 20 minutes.

Strobe® Pro is compatible with many commonly used fungicides, liquid fertilizers, herbicides, insecticides and biological control products. If tank mixes are desired, observe all directions, precautions and limitations on labeling of all products used.

Consult compatibility charts or your local or State turf authority for compatibility information. Do not combine Strobe® Pro in the spray tank with pesticides, surfactants, or fertilizers, unless compatibility charts or your own prior use has shown that the combination is physically compatible, effective, and non-injurious under your conditions of use. If physical compatibility is unknown, the following procedure should be followed. Pour the specified proportions of the products into a suitable container of water, mix thoroughly and allow to stand at least twenty (20) minutes, if the combination remains mixed or can be re-mixed readily, the mixture is considered physically compatible.

### **SPECIFIC USE DIRECTIONS**

Strobe® Pro is specified for control of certain pathogens causing foliar, stem, and root diseases, including leaf and stem blights, leaf spots, patch diseases, mildews, anthracnose, fairy rings, molds, and rusts of turfgrass plants. Strobe® Pro may be used to control certain diseases on turf in golf courses, lawns and landscape areas around residential, institutional, public, commercial and industrial buildings, parks, recreational areas and athletic fields.

Integrated Pest (Disease) Management (IPM): Sound turf management resulting in healthy, vigorous turf is the foundation of a good IPM program. Cultural practices such as proper choice of turf variety, nutrient management, proper cutting height, thatch management, and proper watering, drainage, and moisture stress management should be integrated with the use of fungicides to increase turf vigor and reduce the susceptibility to disease. Immunoassay detection kits and extension service diagnostic services can assist in the early and accurate identification of causal organisms and corresponding selection of the proper fungicide when required.

**Application Directions:** Strobe® Pro should be applied prior to disease development. Mix Strobe® Pro with the required amount of water and apply as a dilute spray application in 2-4 gals. of water per 1000 sq. ft. (87-174 gals./A).

Apply Strobe® Pro at rates and timings as described in this label.

Repeat applications at specified intervals for as long as required. For spot treatments, use 1.5 fl. oz. Strobe® Pro per 1 to 2 gal. of water.

### **Restrictions**

Do not apply more than 8.1 gal. product/acre/year (23.75 fl. oz. product/1000 sq. ft./year)

Do not apply more than 7.047 lbs. a.i. propiconazole/acre/year.

Do not apply more than 4.21 lbs. a.i. azoxystrobin/acre/year.

Do not apply more than 32 times a year at the 0.75 oz. rate/1000 sq. ft.

Do not apply more than 16 times a year at the 1.5 oz. rate/1000 sq. ft.

Do not apply more than 8 times a year at the 3.0 oz. rate/1000 sq. ft.

Applications may be made by ground only.

**For use with soil injection applications:** Strobe® Pro may be applied through a liquid fungicide injector for the

control of ectotrophic root diseases such as summer patch and take-all patch. Use Strobe® Pro only in liquid injection equipment specifically designated for pesticide use.

Apply Strobe® Pro at 1.5-3 fl. oz. per 1000 sq. ft. Spray carrier volume must fall within 30-150 gals, of water per 1000 sq. ft. Injection hole spacing of 1 inch by 1 inch is specified for optimum control. Injection depth must be no greater than 2 inches. One inch depth is specified for optimum results. Application must follow disease control strategies used

for normal broadcast spray programs.

**For use in the establishment of turfgrass from seed or in overseeding of dormant turfgrass:** Strobe® Pro may be used for control of certain turfgrass diseases associated with turfgrass establishment from seed. Strobe® Pro may also be used during overseeding of dormant turfgrass.

Strobe® Pro may be safely applied before or after seeding or at seedling germination and emergence to ryegrass, bentgrass, bluegrass, and fescue turfgrass types. Optimum application timing is during seeding.

**See Application Directions section.**

**Rate Ranges:** Use the shorter specified application interval and/or use the higher specified rate when prolonged favorable disease conditions exist.



Target Diseases	Use Rate (fl. oz. product per 1000 sq. ft.)	Application Interval (Days)	Application Instructions*
Anthracnose ( <i>Colletotrichum cereale</i> ) (formerly known as <i>C. graminicola</i> )	1.5 – 3.0	14 – 28	Use preventatively. Begin applications when conditions are favorable for disease infection, prior to disease symptom development.  The maximum application interval for greens is 14 days.
Bentgrass Dead Spot ( <i>Ophiosphaerella Agrostis</i> )	1.5 – 3.0	14	Begin applications prior to disease development.
Brown Patch ( <i>Rhizoctonia solani</i> )	0.75 – 1.5	14	For higher cut cool season turfgrass, St. Augustine grass and centipede grass (fairways, lawns, sodfarms, etc, above 0.375 inch in height). Apply when conditions are favorable for disease development.
Brown Patch ( <i>Rhizoctonia solani</i> ) and other Rhizoctonia ( <i>R. spp.</i> ) diseases such as Large Patch, Zoysia Patch, Leaf & Sheath Spot ( <i>R. zeae</i> )	1.5 – 3.0	14 – 28	For Brown Patch, apply when conditions are favorable for disease development.  For Large Patch of all warm season turfgrasses, make 1 or 2 applications in the fall prior to infection or when conditions are favorable for infection.  For Zoysia Patch, make 1 or 2 applications approximately one month prior to Zoysia grass dormancy. Reapply 14 to 28 days later.  For Leaf & Sheath Spot ( <i>R. zeae</i> ), apply when conditions are favorable for infection such as sequential or periods of temperatures at or above 90 degrees Fahrenheit. Curative control may necessitate several applications. Target spray applications at crown of turfgrass.

Continued

Target Diseases	Use Rate (fl. oz. product per 1000 sq. ft.)	Application Interval (Days)	Remarks*
Cool Weather Brown Patch, Yellow Patch ( <i>Rhizoctonia cereale</i> )	3.0	28	Make 1 or 2 applications in fall or when conditions are favorable for disease development.
Dichondra Rust ( <i>Puccinia dichondrae</i> )	3.0	14 – 21	Apply when conditions are favorable for disease development.
Dollar Spot ( <i>Sclerotinia homoeocarpa</i> )	0.75 – 1.5	14	Apply preventively on higher cut cool season turfgrass (fairways, lawns, etc. mowed at or above 0.375 inch in height during periods of low to moderate disease pressure).
Dollar Spot ( <i>Sclerotinia homoeocarpa</i> )	1.5 – 3.0	14 – 28	Apply preventively when conditions are favorable for disease development. For non-residential turf with active dollar spot, use the higher rate and combine with Chlorothalonil.
Fairy Ring ( <i>Lycoperdon spp.</i> , <i>Agrocybe pediades</i> , and <i>Bovistia plumbea</i> )	1.5 – 3.0	14 – 28	Apply as soon as possible after fairy ring symptoms develop. Apply in 2-4 gal. of water per 1000 sq. ft. Add the specified rate of a wetting agent to the final spray. Severely damaged or thin turf may require reseeding. Fairy ring symptoms may take 2 to 3 weeks to disappear following application. If area is hydrophobic, use wetting agents and irrigate prior to application(s) of Strobe® Pro. Reapplication after 28 days may be required in some cases.
Gray Leaf Spot ( <i>Pyricularia grisea</i> )	1.5 – 3.0	14 – 28	Use Strobe® Pro in a preventive disease control program. Begin applications before disease is present and alternate with other fungicide chemistries that control gray leaf spot.

Continued

Target Diseases	Use Rate (fl. oz. product per 1000 sq. ft.)	Application Interval (Days)	Remarks*
Gray Snow Mold Typhula Blight ( <i>Typhufa incarnata</i> )	3.0	14 – 28	Make two applications of 3 fl. oz. spaced 14-28 days apart in late fall just before snow cover. Tank mixing with another snow mold fungicide, such as Chlorothalonil and Fludioxonil, may enhance control under severe disease pressure.
Leaf Rust Stem Rust Stripe Rust ( <i>Puccinia spp.</i> )	1.5 – 3.0	14 – 28	Begin applications when conditions are favorable for disease infection, prior to disease symptom development.
Leaf Spot ( <i>Bipolaris sorokiniana</i> )	1.5 – 3.0	14 – 21	Apply when conditions are favorable for disease development.
Melting Out ( <i>Drechslera poae</i> )	1.5 – 3.0	14 – 21	Apply when conditions are favorable for disease development.
Microdochium Patch (formerly known as Fusarium Patch ( <i>Microdochium nivale</i> ))	1.5 – 3.0	14 – 28	Use preventatively. Begin applications when conditions are favorable for disease infection, prior to disease symptom development.
Necrotic Ring Spot ( <i>Ophiosphaerella korrae</i> )	3.0	14 – 28	Apply when conditions are favorable for disease development.
Pink Patch ( <i>Limonomyses roseipellis</i> )	1.5 – 3.0	14 – 28	Apply when conditions are favorable for disease development.
Pink Snow Mold ( <i>Microdochium nivale</i> )	3.0	10 – 28	Make two applications of 3 fl. oz. spaced 10-28 days apart in late fall just before snow cover. Tank mixing with another snow mold fungicide, such as Chlorothalonil and Fludioxonil, may enhance control under severe disease pressure.

Continued

Target Diseases	Use Rate (fl. oz. product per 1000 sq. ft.)	Application Interval (Days)	Remarks*
Powdery Mildew ( <i>Erysiphe graminis</i> )	1.5 – 3.0	14 – 28	Begin applications when conditions are favorable for disease infection, prior to disease symptom development.
Pythium Blight Pythium Root Rot ( <i>Pythium aphanidermatum</i> , <i>Pythium spp.</i> )	3.0	14	Use preventatively. Begin applications when conditions are favorable for disease infection prior to disease symptom development. During periods of prolonged favorable conditions, treat on the 10-day application interval. For use on newly seeded as well as established turf.
Red Thread ( <i>Laetisaria fuciformis</i> )	1.5 – 3.0	14 – 28	Apply when conditions are favorable for disease development.
Southern Blight ( <i>Scierotium rolfsii</i> )	1.5 – 3.0	14 – 28	Apply when conditions are favorable for disease development.
Spring Dead Spot ( <i>Ophiosphaerella korrae</i> ) or ( <i>Ophiosphaerella narmari</i> ) or ( <i>Ophiosphaerella herpotricha</i> )	3.0	14 – 28	Apply 1 or 2 applications approximately one month prior to bermudagrass dormancy. 1/4" to 1/2" of irrigation directly after application is required. Reapply 14 to 28 days later.
Summer Patch ( <i>Magnaporthe poae</i> )	1.5 – 3.0	14 – 28	Apply when conditions are favorable for disease development. Initiate applications when soil temperatures reach 65°F at a 2-inch soil depth.
Take-all Patch ( <i>Gaeumannomyces gram inisvan a venae</i> )	3.0	28	Begin applications when conditions are favorable for disease infection prior to disease symptom development. Make two applications, 28 days apart in the spring and two applications 28 days apart in the fall.

\*Do not apply more than two sequential applications of Strobe® Pro for control of Gray Leaf Spot and Pythium spp. For all other diseases when Gray Leaf Spot and Pythium spp. are not present, do not apply more than three sequential applications of Strobe® Pro.

**Strobe® Pro Rate Conversion Chart for Turf**

<b>Fl. Oz. Product* Per 1000 Sq. Ft.</b>	<b>Pints Product Per Acre</b>	<b>Gallons of Product Per Acre</b>
0.75	2.0	0.26
1.0	2.7	0.34
1.5	4.0	0.51
2.0	5.4	0.68
2.5	6.8	0.85
3.0	8.2	1.02

\*One fluid ounce of Strobe® Pro contains 0.1088 ounces (weight) propiconazole and 0.065 ounces (weight) azoxystrobin.

Amount of Strobe® Pro for Turf Applications per 100 Gallons

**Spray volume: 2 gals./1000 sq. ft.**

<b>Strobe® Pro Use Rate (fl. oz./1000 sq. ft.)</b>	<b>Pints of product/100 gallons</b>
0.75	2.3
1.0	3.1
1.5	4.7
2.0	6.2
2.5	7.8
3.0	9.4

**Spray volume: 3 gals./1000 sq. ft.**

<b>Strobe® Pro Use Rate (fl. oz./1000 sq. ft.)</b>	<b>Pints of product/100 gallons</b>
0.75	1.5
1.0	2.1
1.5	3.0
2.0	4.1
2.5	5.2
3.0	6.2

Spray volume: 4 gals./1000 sq. ft.

Strobe® Pro Use Rate (fl. oz./1000 sq. ft.)	Pints of product/100 gallons
0.75	1.15
1.0	1.55
1.5	2.25
2.0	3.1
2.5	3.9
3.0	4.65

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in original containers only. Keep container closed when not in use. Do not store near food or feed. In case of spill on floor or paved surfaces, mop up and remove to chemical waste storage area until proper disposal can be made if product cannot be reused according to the label.

**PESTICIDE DISPOSAL:** Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative of the nearest EPA Regional Office for guidance.

**CONTAINER HANDLING:** Non-refillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

## LIMITATION OF WARRANTY AND LIABILITY

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Control Solutions, Inc. or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Control Solutions, Inc. and Seller harmless for any claims relating to such factors.

Control Solutions, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or Control Solutions, Inc., and Buyer and User assume the risk of any such use. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW CONTROL SOLUTIONS, INC. MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.**

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW IT IS CONTROL SOLUTIONS INC., AND SELLERS INTENTION THAT** in no event shall Control Solutions or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF CONTROL SOLUTIONS, INC., AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF CONTROL SOLUTIONS, INC. OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

Control Solutions, Inc. and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Control Solutions, Inc.

Quali-Pro® is a registered trademark of Control Solutions, Inc.

Strobe® is a registered trademark of Control Solutions, Inc.







GROUP 3 11 FUNGICIDE



# STROBE® PRO

BROAD SPECTRUM FUNGICIDE

For control of Turf Diseases.

ACTIVE INGREDIENTS:	% BY WT.
Azoxystrobin* .....	6.0%
Propiconazole** .....	10.0%
OTHER INGREDIENTS: .....	84.0%
TOTAL: .....	100.0%

KEEP OUT OF REACH OF CHILDREN  
**CAUTION**

EPA Reg. No: 53883-418

EPA Est No. 53883-TX-002

A liquid soluble concentrate that contains 0.87 lbs. propiconazole and 0.52 lbs. azoxystrobin per gallon.

\*IUPAC: methyl (E)-2-[2-[6-(2-cyanophenoxy)pyrimidin-4-yloxy]phenyl]-3-methoxyacrylate

\*\*CAS Number 60207-90-1

## CONTENTS: 1 GALLON



Manufactured for:  
**Control Solutions Inc.**

5903 Genoa-Red Bluff, Pasadena, TX 77507  
A member of Adama Consumer and Professional Solutions

PEEL BACK HERE